

# PMIP4 HadGEM2-A dust data

The simulation of the pre-industrial and LGM climate states with HadGEM2-A are described in more detail by Hopcroft & Valdes (2015), whilst the dust results are described by Hopcroft et al (2015). The dust scheme itself has been documented by Woodward (2011). HadGEM2-ES is documented by Collins et al (2011).

## Grid

## Time steps and data files

- files names = PMIP4\_DUST\_Hopcroft\_<field>\_<expt>.nc
  - field = see [Variables](#) section below
  - expt = PI or LGM

## Variables

### Notes

- The dust size bin edges are: 0.0316, 0.1, 0.316, 1.0, 3.16, 10.0 and 31.6  $\mu\text{m}$ .
- AOD is given on 6 wavelengths. These are 0.38, 0.44, 0.55, 0.67, 0.87 and 1.02  $\mu\text{m}$  (Bellouin et al 2007).

### 2D dust fields

- **emis**: emission (lon,lat,size,month) [g/m<sup>2</sup>/a]
- **ddep**: dry deposition (lon,lat,size,month) [g/m<sup>2</sup>/a]
- **wdep**: wet deposition (lon,lat,size,month) [g/m<sup>2</sup>/a]
- **load**: dust loading (lon,lat,size,month) [kg/m<sup>2</sup>]
- **aot**: aerosol optical thickness (lon,lat,wavelength,month) [unitless]
- **rfss**: surface short-wave radiative forcing (lon,lat,month) [W/m<sup>2</sup>]
- **rfsl**: surface long-wave radiative forcing (lon,lat,month) [W/m<sup>2</sup>]
- **rfts**: TOA short-wave radiative forcing (lon,lat,month) [W/m<sup>2</sup>]
- **rftl**: TOA long-wave radiative forcing (lon,lat,month) [W/m<sup>2</sup>]

### 3D dust fields

- **conc**: mass mixing ratio (lon,lat,lev,month) [kg/kg] x 6 bins

## References

- Bellouin, N et al., (2007), **Improved representation of aerosols for HadGEM2**, [Hadley Centre Technical Note 73](#), Met Office Hadley Centre, Exeter, UK:
- Collins, WJ et al (2011), **Development and evaluation of an Earth-System model - HadGEM2**, Geoscientific Model Development, 4, 1051-1075, doi:[10.5194/gmd-4-1051-2011](https://doi.org/10.5194/gmd-4-1051-2011).
- Hopcroft, P.O. and P.J. Valdes (2014), **Last Glacial Maximum constraints on the Earth System Model HadGEM2-ES**, Climate Dynamics, 45(5), 1657-1672, doi:[10.1007/s00382-014-2421-0](https://doi.org/10.1007/s00382-014-2421-0).
- Hopcroft, P.O., P.J. Valdes, S. Woodward and M. Joshi (2015), **Last glacial maximum radiative forcing from mineral dust aerosols in an Earth System model**, J Geophysical Research, 120(16), 8186-8205, doi:[10.1002/2015JD023742](https://doi.org/10.1002/2015JD023742).
- Woodward, S. (2011), **Mineral dust in HadGEM2**, [Hadley Centre Technical Note 87](#), Met Office Hadley Centre, Exeter, UK.

## Download

From:

<https://pmip4.lsce.ipsl.fr/> - **PMIP4**

Permanent link:

[https://pmip4.lsce.ipsl.fr/doku.php/data:dust\\_hadgem2\\_a?rev=1469439900](https://pmip4.lsce.ipsl.fr/doku.php/data:dust_hadgem2_a?rev=1469439900)

Last update: **2016/07/25 11:45**

