

Design for the Last Deglaciation experiment

You will find on this page information about the experiment design for the PMIP4 [Last Deglaciation](#) experiments.

Please make sure to read the [Associated publications](#) before setting up your experiments or using the output data, and read any *how-to* sections associated with specific boundary conditions.



Get in touch with the following people if you have questions:

| | |
|--|-------------------------------------|
| Jean-Yves Peterschmitt | Technical questions or missing data |
| Ruza Ivanovic | working group leader |
| Lauren Gregoire | working group co-leader |
| Masa Kageyama | working group steering |
| Didier Roche | working group steering |
| Paul Valdes | working group steering |
| Andrea Burke | working group steering |

Associated publications

Last deglaciation experiment design, version 1:

Ivanovic, R. F.; Gregoire, L. J.; Kageyama, M.; Roche, D. M.; Valdes, P. J.; Burke, A.; Drummond, R.; Peltier, W. R. and Tarasov, L.: **Transient climate simulations of the deglaciation 21-9 thousand years before present, version 1; PMIP4 Core experiment design and boundary conditions**, Geosci. Model Dev., 9, 2563–2587, doi:[10.5194/gmd-9-2563-2016](https://doi.org/10.5194/gmd-9-2563-2016), 2016.

Version 1 Specifications

For general advice on boundary condition implementation in palaeoclimate models, see [Kageyama et al. \(2016\)](#).

Last Glacial Maximum spinup (21 ka)

This spinup simulation is compatible with the PMIP4-CMIP6 LGM experiment, which can also be used as the initialisation state for the fully transient run from 21 ka onwards, provided the ICE-6G_C or GLAC-1D ice sheet reconstructions and associated boundary conditions (orography, coastlines and bathymetry) were used.

| | |
|------------|-----------------------------|
| | PMIP4 specifications |
| PMIP4 name | LDv1-LGMspin |

| | PMIP4 specifications |
|--------------------------------------|--|
| Astronomical parameters | eccentricity = 0.018994 obliquity = 22.949° perihelion-180° = 114.42° Date of vernal equinox : Noon, 21st March |
| Solar constant | 1361.0 ± 0.51365 W m ⁻² |
| Trace gases | CO₂ = 190 ppm CH₄ = 375 ppb N₂O = 200 ppb CFC = 0 O₃ = Preindustrial (e.g. 10 DU) |
| Ice sheets, orography and coastlines | 21 ka data from either - ICE-6G_C reconstruction: [Access to data] - GLAC-1D reconstruction: [Access to data] |
| Bathymetry | Keep consistent with the coastlines, using either: - Data associated with the ice sheet - Preindustrial bathymetry |
| Global ocean salinity | + 1 psu, relative to preindustrial |
| All others | As per the PMIP4-CMIP6 LGM experiment |

Transient orbit and trace gases spinup (26-21 ka)

This option for spinning-up the last deglaciation simulation uses transient orbital parameters and trace gases from 26-21 ka.

| | PMIP4 specifications |
|-------------------------|--|
| PMIP4 name | LDv1-transpin |
| Astronomical parameters | All orbital parameters should be transient as per Berger (1978) 26-21 ka [Access to data & README !] |
| Trace gases | All adjusted to the AICC2012 chronology Veres et al. (2013) 26-21 ka: CO₂ = Transient, as per Bereiter et al. (2015) : [Access to data (md5sum = c54a033d8cbf588bc2b95d3b92ff9b1c)] CH₄ = Transient, as per Louergue et al. (2008) : [Access to data] N₂O = Transient, as per Schilt et al. (2010) : [Access to data] |
| <i>All others</i> | As per the LGM (21 ka) spinup type; LDv1-LGMspin |

Transient deglaciation (21-0 ka)

These are the specifications for the full transient run 21-0 ka.
(Note, the period of focus for version 1 of the experiment is 21-9 ka, but all boundary conditions are provided until 0 ka so that groups can extend the run to present if they wish).

| | PMIP4 specifications |
|--------------------------------|--|
| PMIP4 name | LDv1 |
| Initial conditions (pre 21 ka) | Recommended (optional) to use either: - LDv1-LGMspin - LDv1-transpin See above for details. The method must be documented, including information on the state of spinup |

| | PMIP4 specifications |
|--|--|
| Astronomical parameters | Transient, as per Berger (1978) [Access to data & README !] |
| Solar constant | 1361.0 ± 0.51365 W m ⁻² |
| Trace gases | Adjusted to the AICC2012 chronology Veres et al. (2013) 21-0 ka: CO₂ = Transient, as per Bereiter et al. (2015) : [Access to data (md5sum = c54a033d8cbf588bc2b95d3b92ff9b1c)] CH₄ = Transient, as per Louergue et al. (2008) : [Access to data] N₂O = Transient, as per Schilt et al. (2010) : [Access to data] CFC = 0 O₃ = Preindustrial (e.g. 10 DU) |
| Ice sheet | Transient, with a choice of either : - ICE-6G_C reconstruction: [Access to data] - GLAC-1D reconstruction: [Access to data] How often to update the ice sheet is optional |
| Orography and coastlines | Transient. To be consistent with the choice of ice sheet. Orography is updated on the same timestep as the ice sheet. It is optional how often the land-sea mask is updated, but ensure consistency with the ice sheet reconstruction is maintained |
| Bathymetry | Keep consistent with the coastlines, and otherwise use either: - Data associated with the ice sheet; it is optional how often the bathymetry is updated - Preindustrial bathymetry |
| River routing | Ensure that rivers reach the coastline It is recommended (optional) to use one of the following: - Preindustrial configuration for the model - Transient routing provided with the ice sheet reconstruction (if available) - Manual/model calculation of river network to match topography |
| Freshwater fluxes | At participant discretion. Three options are: <i>melt-uniform</i> , <i>melt-routed</i> and <i>no-melt</i> : - <i>Melt-uniform</i> : use a globally uniform ice meltwater flux, e.g. as associated with one of the ice sheet reconstructions [ICE-6G_C] - [GLAC-1D] - <i>Melt-routed</i> : use a routed ice meltwater flux, e.g. as associated with one of the ice sheet reconstructions [ICE-6G_C] - [GLAC-1D] - <i>No-melt</i> : have no ice sheet meltwater in the simulation It is recommended (optional) to run at least one Core simulation with a scenario consistent with the chosen ice sheet reconstruction to conserve salinity. |
| Vegetation & land cover Aerosols (dust) | Prescribed preindustrial cover or dynamic vegetation model Prescribed preindustrial distribution or prognostic aerosols |

Focused simulations

- Empty
- Empty

References cited

- Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S.

- and Chappellaz, J.: Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present, *Geophys. Res. Lett.*, 42(2), 2014GL061957, [doi:10.1002/2014GL061957](https://doi.org/10.1002/2014GL061957), 2015.
- Berger, A.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, *J. Atmospheric Sci.*, 35(12), 2362–2367, [doi:10.1175/1520-0469\(1978\)035<2362:LTVODI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2), 1978.
 - Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclauss, J., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q. and Zhou, T.: PMIP4-CMIP6: the contribution of the Paleoclimate Modelling Intercomparison Project to CMIP6, *Geosci. Model Dev. Discuss.*, 1–46, [doi:10.5194/gmd-2016-106](https://doi.org/10.5194/gmd-2016-106), 2016.
 - Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F. and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453(7193), 383–386, [doi:10.1038/nature06950](https://doi.org/10.1038/nature06950), 2008.
 - Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J., Loulergue, L., Schüpbach, S., Spahni, R., Fischer, H. and Stocker, T. F.: Atmospheric nitrous oxide during the last 140,000 years, *Earth Planet. Sci. Lett.*, 300(1–2), 33–43, [doi:10.1016/j.epsl.2010.09.027](https://doi.org/10.1016/j.epsl.2010.09.027), 2010.
 - Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim Past*, 9(4), 1733–1748, [doi:10.5194/cp-9-1733-2013](https://doi.org/10.5194/cp-9-1733-2013), 2013.

From:

<https://pmip4.lsce.ipsl.fr/> - **PMIP4**

Permanent link:

https://pmip4.lsce.ipsl.fr/doku.php/exp_design:degla?rev=1470758158

Last update: **2016/08/09 15:55**

