

Design for the Penultimate Deglaciation experiment

You will find on this page information about the experiment design for the PMIP4 [Penultimate Deglaciation](#) experiments.

Please make sure to read the [Associated publications](#) before setting up your experiments or using the output data, and read any *how-to* sections associated with specific boundary conditions.



Get in touch with the following people if you have questions:

Laurie Menviel	Experimental design questions
Emilie Capron	Experimental design questions
Ruza Ivanovic	working group leader
Jean-Yves Peterschmitt	Technical questions or missing data

Associated publications

- **Penultimate deglaciation experiment design, version 1:**

The penultimate deglaciation: protocol for PMIP4 transient numerical simulations between 140 and 127 ka, version 1.0, Menviel et al, GMD, 2019,
https://dx.doi.org/available_soon

Version 1 Specifications

For general advice on boundary condition implementation in palaeoclimate models, see [Kageyama et al. \(2016\)](#).

Penultimate Glacial Maximum spinup (140 ka)

If possible, this spinup simulation should start from the PMIP4-CMIP6 LGM (21 ka) experiment, as equilibrium would be reached more quickly.

	PMIP4 specifications
PMIP4 name	PDv1-LGMspin
Astronomical parameters	eccentricity = 0.033 obliquity = 23.42° perihelion-180° = 251° Date of vernal equinox : Noon, 21st March
Solar constant	1361.0 ± 0.51365 W m ⁻²

	PMIP4 specifications
Trace gases	CO₂ = 195 ppm CH₄ = 387 ppb N₂O = 201 ppb CFC = 0 O₃ = Preindustrial (e.g. 10 DU)
Ice sheets, orography and coastlines	140 ka data from Combined ice-sheet reconstruction (IcIES-NH, GSM-G and GSM-A): [Access to data]
Bathymetry	Keep consistent with the coastlines, using either: - Data associated with the ice sheet - Preindustrial bathymetry
Global ocean salinity	+ 0.85 psu, relative to preindustrial
All others	See manuscript section 6.1

Transient penultimate deglaciation (140-127 ka)

These are the specifications for the full transient run 140-127 ka.

	PMIP4 specifications
PMIP4 name	PDv1
Initial conditions (pre 21 ka)	Recommended: PDv1-LGMspin See above for details. The method must be documented, including information on the state of spinup
Astronomical parameters	Transient, as per Berger (1978) [Access to data & README !]
Solar constant	$1361.0 \pm 0.51365 \text{ W m}^{-2}$
Trace gases	CO₂ = Transient, as per the spline of Koehler et al. (2017) : [Access to data] CH₄ = Transient, as per the spline of Koehler et al. (2017) : [Access to data] N₂O = Linear increase from 201 ppb at 140 ka to 218.74 ppb at 134.5 ka then transient, as per the spline of Koehler et al. (2017) : [Access to data] CFC = 0 O₃ = Preindustrial (e.g. 10 DU)
Ice sheet	Transient: Combined ice-sheet reconstruction (IcIES-NH, GSM-G and GSM-A) [Access to data] How often to update the ice sheet is optional
Orography and coastlines	Transient. To be consistent with the choice of ice sheet. Orography is updated on the same timestep as the ice sheet. It is optional how often the land-sea mask is updated, but ensure consistency with the ice sheet reconstruction is maintained
Bathymetry	Keep consistent with the coastlines, and otherwise use either: - Data associated with the ice sheet; it is optional how often the bathymetry is updated - Preindustrial bathymetry
River routing	Ensure that rivers reach the coastline It is recommended (optional) to use one of the following: - Self-consistent paleo-routing described in section 6.2.3 - Preindustrial configuration for the model - Manual/model calculation of river network to match topography

	PMIP4 specifications
Freshwater fluxes	Recommended North Atlantic option is <i>fSL</i> and a constant 0.0135 Sv flux around the Antarctic coast between 140-130 ka - <i>fSL</i> : meltwater flux based on changes in sea-level [] - [] - <i>fIRD</i> : meltwater flux based on Norwegian Sea and North Atlantic IRD [] - [] - <i>fIC</i> : meltwater flux based on ice-sheet changes [] - [] - <i>fSL2</i> : meltwater flux based on changes in sea-level and triangular input max. 0.15 Sv between 131-128 ka on the Antarctic coast [] - [] - <i>fUN</i> : Globally uniform meltwater input based on sea-level changes
Vegetation & land cover Aerosols (dust)	Prescribed preindustrial cover or dynamic vegetation model Prescribed preindustrial distribution or prognostic aerosols

Focused simulations

- Empty
- Empty

Paleorecords to use for model-data comparisons

Overview

See Table 3 and Table 4 of the [Penultimate Deglaciation GMD paper](#)

Table 3									Table 4									
Click on a table to get a bigger version, or download the GMD paper																		
Tracer interpretation	Core	Coordinates and depth (m)	σ^1 (ka)	σ^2 (ka)	σ^3 (ka)	σ^4 (ka)	σ^5 (ka)	References	Tracer interpretation	Core, coordinates and depth/elevation	Chronology	σ^1 (ka)	σ^2 (ka)	σ^3 (ka)	σ^4 (ka)	σ^5 (ka)	References	
Sea-level									Atm. CO ₂ concentration									
Sea-level	Red Sea cores	-	137.0±0.7 increases		133.4±0.7 main increase		130.2±1	Grant et al. (2012) This study	Atm. CO ₂ concentration	EDC	75.05°S, 123.19°E, 3233 m asl	Ice core	137.8±2.7 increases				128.0±1.8	Beer et al. (2015)
Benthic $\delta^{18}O$									SST									
North Atlantic intermediate-depth ventilation	ODP983	60.40°N, 23.64°W, 1984 m	136.1±1.2 weaker vent				128.1±0.9	Raymo et al. (2004) Barber et al. (2015)	North Atlantic Summer SST (FFA)	ODP980	55.80°N, 14.11°W, 2180 m depth	Alignment onto Corecha U-Th based chrono.				128.7±1.3	128.7±1.3	Oppo et al. (2006)
North Atlantic deep-water ventilation	ODP980	55.80°N, 14.11°W, 2180 m	137±1.9			128.6±1.8	127.6±1.3	Oppo et al. (2006)	North Atlantic Summer SST (FFA)	SL90-03	40.51°N, 32.05°W, 2475 m depth	Alignment onto Corecha U-Th based chrono.	136.9±1.6 colder [E]			131±1	CS98	Cortijo et al. (1999)
North Atlantic deep-water ventilation	MD95-2042	37.80°N, 10.17°E, 3140 m				131.0±1.4 stronger vent (T)		Shackleton et al. (2003)	W. Meditter. Sea SST (IS'37)	ODP 976	36.20°N, 4.31°E, 1108 m depth	Alignment onto Corecha U-Th based chrono.	135.9±1.1 colder [A]	134 ±1.1 [B]	133.3** ±0.9 [C]	131.4±0.9 warmer [D]	Martini et al. (2014)	
Southern Ocean deep-water ventilation	Stack of UI308 CH169-K09 and ODP 1063	49.88°N, 24.24°W, 3883 m 41.36°N, 47.35°W, 4100 m 33.60°N, 57.62°W, 4584 m	135.9±2.0			130.3±1.6	129.2±1.4	Hubert et al. (1999) Lajoye et al. (2000) Deane et al. (2017)	Southern Ocean Summer SST (FFA)	MD02-2488	46.48°S, 88.02°E, 3420 m depth	Alignment onto AICC2012				130±2.1** max. warm [B]	Gavin et al. (2009) Gavin et al. (2012)	
Southern Ocean deep-water ventilation	MD02-2488	46.48°S, 88.02°E, 3420 m				131.9±2.1 stronger vent (U)	130.2±2.2 weaker vent (V)	Gavin et al. (2012)	Southern Ocean SST (Mg/Ca)	MD02-2120	45.54°S, 174.02°E, 1210 m depth	Alignment onto AICC2012				128.1±2.5** max. warm [S]	Phillips et al. (2005)	
Planktic $\delta^{18}O$ and $\delta^{17}O$ sw									Air temperature (SAT)									
North Atlantic surface $\delta^{18}O$ salinity	ODP980	55.80°N, 14.11°W, 2180 m				130.0±1.3		Oppo et al. (2006)	Antarctic Annual SAT (ice d0)	EDC ice core	75.1°S, (23.55°E), 3233 m a.s.l.	Ice core	135.6±2.5 (warmer) [W]			129.4±1.8** max. warm [X]	Jouzel et al. (2007)	
North Atlantic surface $\delta^{18}O$ salinity	SL90-03	40.51°N, 32.05°W				131.0±1.1		CS98	SAT	Corecha Cave, Italy	43.97°N, 13.07°E, 840 m a.s.l.	Alignment onto Corecha U-Th based chrono.				131.0±0.7 warmer [K]	128.6±0.6 warm plateau [L]	Drysdale et al., 2010
North Atlantic surface $\delta^{18}O$ salinity	MD95-2042	37.80°N, 10.17°E				131.6±1.5		Marino et al. (2013)	Vegetation/precipitation in Southern Europe	Lago di Monticchio, Italy	40.95°N, 15.58°E, 656 m a.s.l.	Independent absolute varve chronology				127.2±1.6 max. warm/cool [G]	Brauer et al. (2007)	
North Atlantic surface $\delta^{18}O$ salinity	ODP 976	36.20°N, 4.31°E				133.9±0.9		Martini et al. (2014)	(Temperate tree pollen)	Isosmia tere sequences, Greece	39.45°N, 20.01°E, 470 m a.s.l.	Orbital tuning				132.7±2.3 water [R]	Tzedakis et al. (2003)	
Speleothem $\delta^{18}O$									Precipitation									
North Atlantic surface $\delta^{18}O$	Corecha Cave, Italy	43.97°N, 13.07°E, 840 m a.s.l.	133.9±1.2 A1 meltwater input (D)			131.0±0.7 NA meltwater pinched (J)		Drysdale et al. (2010) Tzedakis et al. (2018) Martini et al. (2015)	Intensity of Asian monsoon ($\delta^{18}O$)	Chinese Caves	25.28°N to 32.5°N, 108.08-119.16°E, 080 - 1900 m a.s.l.	Absolute U-Th based chronology	135.6±0.5 drier [M]	133.7 ±0.5 [N]	133.3** ±0.5 [O]	128.9±0.1 water [P]	Cheng et al. (2016)	
Mean ages for the beginning of σ^1 - σ^5 from Tables 3 and 4									Mean ages for the beginning of σ^1 - σ^5 from Tables 3 and 4									
Mean ages			136.4±1.7	133.9±0.8	133.3±1.1	130.4±1.3	128.5±1.3		Intensity of East Asian monsoon (grain size)	Chinese loess	35.62-37.14°N, 103.20 - 109.85°E	Alignment onto Holo-Sarhao U-Th based chrono.				130.2±0.4 water [Q]	Yang and Ding (2014)	
Table 3									Table 4									

Data

CH69-K09
MD95-2042
ODP976
ODP980
ODP983
ODP1063

[SU90-03](#)

References cited

- Berger, A.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, *J. Atmospheric Sci.*, 35(12), 2362–2367, [doi:10.1175/1520-0469\(1978\)035<2362:LTVODI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2), 1978.
- Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclauss, J., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q. and Zhou, T.: PMIP4-CMIP6: the contribution of the Paleoclimate Modelling Intercomparison Project to CMIP6, *Geosci. Model Dev. Discuss.*, 1–46, [doi:10.5194/gmd-2016-106](https://doi.org/10.5194/gmd-2016-106), 2016.
- P. Koehler and C. Nehrbass-Ahles and J. Schmitt and T.F. Stocker and H. Fischer: A 156 kyr smoothed history of the atmospheric greenhouse gases CO₂, CH₄, and N₂O and their radiative forcing, *Earth System Science Data*, 9, 363–387, [doi:10.5194/essd-9-363-2017](https://doi.org/10.5194/essd-9-363-2017), 2017.
- Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim Past*, 9(4), 1733–1748, [doi:10.5194/cp-9-1733-2013](https://doi.org/10.5194/cp-9-1733-2013), 2013.

From:

<https://pmip4.lsce.ipsl.fr/> - **PMIP4**

Permanent link:

https://pmip4.lsce.ipsl.fr/doku.php/exp_design:degla_t2?rev=1549034133Last update: **2019/02/01 16:15**