

Design for the Penultimate Deglaciation experiment

You will find on this page information about the experiment design for the PMIP4 [Penultimate Deglaciation](#) experiments.

This protocol is a product of the *PAGES-PMIP working group on Quaternary Interglacials (QUIGS)*



Please make sure to read the [Associated publications](#) before setting up your experiments or using the output data, and read any *how-to* sections associated with specific boundary conditions.

Get in touch with the following people if you have questions:

Laurie Menviel	Experimental design questions
Emilie Capron	Experimental design questions
Ruza Ivanovic	working group leader
Jean-Yves Peterschmitt	Technical questions or missing data

Associated publications

- **Penultimate Deglaciation experiment design, version 1:**

The penultimate deglaciation: protocol for PMIP4 transient numerical simulations between 140 and 127 ka, version 1.0, Menviel et al, GMD, 2019,
https://dx.doi.org/available_soon

Version 1 Specifications

For general advice on boundary condition implementation in palaeoclimate models, see [Kageyama et al. \(2016\)](#).

Penultimate Glacial Maximum spinup (140 ka)

If possible, this spinup simulation should start from the PMIP4-CMIP6 LGM (21 ka) experiment, as equilibrium would be reached more quickly.

	PMIP4 specifications
PMIP4 name	PDGv1-PGMspin (PDG $\Leftrightarrow$ <i>Penultimate DeGlaciation</i> - PGM $\Leftrightarrow$ <i>Penultimate Glacial Maximum</i>)

	PMIP4 specifications
Astronomical parameters	eccentricity = 0.033 obliquity = 23.414° perihelion-180° = 73° Date of vernal equinox : Noon, 21st March
Solar constant	1361.0 ± 0.51365 W m ⁻²
Trace gases	CO₂ = 191 ppm CH₄ = 385 ppb N₂O = 201 ppb CFC = 0 O₃ = Preindustrial (e.g. 10 DU)
Ice sheets, orography and coastlines	140 ka data from Combined ice-sheet reconstruction (IcIES-NH, GSM-G and GSM-A): [Access to data]
Bathymetry	Keep consistent with the coastlines, using either: - Data associated with the ice sheet - Preindustrial bathymetry
Global ocean salinity	+ 0.85 psu, relative to preindustrial
All others	See manuscript section 6.1

Transient Penultimate Deglaciation (140-127 ka)

These are the specifications for the full transient run 140-127 ka.

	PMIP4 specifications
PMIP4 name	PDGv1
Initial conditions (140 ka)	Recommended: PDGv1-PGMspin See above for details. The method must be documented, including information on the state of spinup
Astronomical parameters	Transient, as per Berger (1978) [Access to data & README !]
Solar constant	1361.0 ± 0.51365 W m ⁻²
Trace gases	CO₂ = Transient, as per the spline of Koehler et al. (2017) : [Access to data] CH₄ = Transient, as per the spline of Koehler et al. (2017) : [Access to data] N₂O = Linear increase from 201 ppb at 140 ka to 218.74 ppb at 134.5 ka then transient, as per the spline of Koehler et al. (2017) : [Access to data] CFC = 0 O₃ = Preindustrial (e.g. 10 DU)
Ice sheet	Transient: Combined ice-sheet reconstruction (IcIES-NH, GSM-G and GSM-A) [Access to data] How often to update the ice sheet is optional
Orography and coastlines	Transient. To be consistent with the choice of ice sheet. Orography is updated on the same timestep as the ice sheet. It is optional how often the land-sea mask is updated, but ensure consistency with the ice sheet reconstruction is maintained
Bathymetry	Keep consistent with the coastlines, and otherwise use either: - Data associated with the ice sheet; it is optional how often the bathymetry is updated - Preindustrial bathymetry

	PMIP4 specifications
River routing	Ensure that rivers reach the coastline It is recommended (optional) to use one of the following: - Self-consistent paleo-routing described in section 6.2.3 - Preindustrial configuration for the model - Manual/model calculation of river network to match topography
Freshwater fluxes	Recommended North Atlantic option is <i>fSL</i> and a constant 0.0135 Sv flux around the Antarctic coast between 140-130 ka [Access to data] - <i>fSL</i> : meltwater flux based on changes in sea-level - <i>fIRD</i> : meltwater flux based on Norwegian Sea and North Atlantic IRD - <i>fIC</i> : meltwater flux based on ice-sheet changes - <i>fSL2</i> : meltwater flux based on changes in sea-level and triangular input max. 0.15 Sv between 131-128 ka on the Antarctic coast - <i>fUN</i> : Globally uniform meltwater input based on sea-level changes
Vegetation & land cover Aerosols (dust)	Prescribed preindustrial cover or dynamic vegetation model Prescribed preindustrial distribution or prognostic aerosols

Focused simulations

- Empty
- Empty

Paleorecords to use for model-data comparisons

Overview

See Table 3 and Table 4 of the [Penultimate Deglaciation GMD paper](#)

Table 3									Table 4									
Click on a table to get a bigger version, or download the GMD paper																		
Tracer interpretation	Core	Coordinates and depth (m)	ψ1 (ka)	ψ2 (ka)	ψ3 (ka)	ψ4 (ka)	ψ5 (ka)	References	Tracer interpretation	Core, coordinates and depth/elevation	Chronology	ψ1 (ka)	ψ2 (ka)	ψ3 (ka)	ψ4 (ka)	ψ5 (ka)	References	
Sea-level									Atm. CO ₂ concentration									
Red Sea cores									EDC: 75.05°S, 123.19°E, 3233 m asl									
Benthic δ ¹⁸ O									Ice core: AICC2012 chrono.									
137.0±0.7 increases									137.8±2.7 increases									
133.4±0.7 main increase									128.0±1.8									
Grant et al. (2012)									Benitez et al. (2015)									
This study																		
Sea-level	Red Sea cores	-	137.0±0.7 increases	133.4±0.7 main increase			130.2±1	Grant et al. (2012)	North Atlantic Summer SST (°F2A)	55.80°N, 14.11°W, 2180 m depth	Alignment onto Corchia U-Th based chrono.					129.7±1.3	128.7±1.3	Oppo et al. (2006)
North Atlantic intermediate-depth ventilation	ODP983	60.40°N, 23.64°W, 1984 m	136.1±1.2 weaker vent				128.1±0.9	Raymo et al. (2004)	North Atlantic Summer SST (°F2A)	40.51°N, 32.05°W, 2475 m depth	Alignment onto Corchia U-Th based chrono.	136.9±1.6 colder [E]				131±1 warmer [F]		CS88 Coriño et al. (1999)
North Atlantic deep-water ventilation	ODP980	55.80°N, 14.11°W, 2180 m	137±1.9				128.6±1.8	127.6±1.3	W. Meditter. Sea SST (Ua-37)	36.20°N, 4.31°E, 1108 m depth	Alignment onto Corchia U-Th based chrono.	135.9±1.1 colder [A]	134 ±1.1 [B]	133.3** ±0.9 [C]		131.4±0.9 warmer [D]		Martini et al. (2014)
North Atlantic deep-water ventilation	MD95-2042	37.80°N, 10.17°E, 3146 m					131.0±1.4 stronger vent (T)	129.2±1.4	Southern Ocean Summer SST (°F2A)	46.48°S, 88.02°E, 3420 m depth	Alignment onto AICC2012					130±2.1** max. warm [K]		Gavin et al. (2009)
North Atlantic deep-water ventilation	Stack of U1308 CH10-K99 and ODP 1063	49.88°N, 24.24°W, 3883 m, 41.76°N, 47.35°W, 4100 m, 33.68°N, 57.62°W, 4584 m	135.9±2.0				130.3±1.6	129.2±1.4	Southern Ocean SST (Mg/Ca)	45.54°S, 114.92°E, 1210 m depth	Alignment onto AICC2012					128.1±2.5** max. warm [S]		Pahske et al. (2003)
South Atlantic deep-water ventilation	MD95-2488	46.48°S, 88.02°E, 3420 m					131.9±2.1 stronger vent (U)	130.2±2.2 weaker vent (V)	Antarctic Annual SAT (ice d0)	75.1°S, 123.35°E, 3233 m a.s.l.	Ice core: AICC2012 chrono.	135.6±2.5 (warmer) [W]				128.4±1.8** max. warm [X]		Jouzel et al. (2007)
South Atlantic deep-water ventilation	MD95-2488	46.48°S, 88.02°E, 3420 m					131.9±2.1 stronger vent (U)	130.2±2.2 weaker vent (V)	SAT	Corchia Cave, Italy	43.97°N, 13.07°E, 840 m a.s.l.	Alignment onto Corchia U-Th based chrono.				131.0±0.7 warmer [K]	128.6±0.6 warm plateau [L]	(Drysdale et al., 2018)
South Atlantic deep-water ventilation	MD95-2488	46.48°S, 88.02°E, 3420 m					131.9±2.1 stronger vent (U)	130.2±2.2 weaker vent (V)	Southern Europe (Mg/L)	Lago di Monticchio, Italy	40.93°N, 15.58°E, 656 m a.s.l.	Independent absolute varve chronology				127.2±1.6 max. warm/cool [G]		Brauer et al. (2007)
North Atlantic surface δ ¹⁸ O	ODP980	55.80°N, 14.11°W, 2180 m					130.0±1.3	129.2±1.4	Vegetation/precipitation in Southern Europe	Lago di Monticchio, Italy	40.93°N, 15.58°E, 656 m a.s.l.	Independent absolute varve chronology				127.2±1.6 max. warm/cool [G]		Brauer et al. (2007)
North Atlantic surface δ ¹⁸ O	SU90-03	40.51°N, 32.05°W, 2180 m					131.0±1.1		(Temperature tree pollen)	Ioannina tree sequence, Greece	39.65°N, 20.91°E, 470 m a.s.l.	Orbital tuning				132.7±2.3 winter [H]		Tzedakis et al. (2003)
North Atlantic surface δ ¹⁸ O	MD95-2042	37.80°N, 10.17°E, 3146 m					133.9±0.9		Intensity of Asian monsoon (δ ¹⁸ O _{ice})	Chinese Caves	25.29°N to 32.5°N, 108.08-119.16°E, 680-1900 m a.s.l.	Absolute U-Th based chronology	135.6±0.5 drier [M]	133.7 ±0.5 [N]	133.3** ±0.5 [O]	128.9±0.1 winter [P]		Cheng et al. (2016)
North Atlantic surface δ ¹⁸ O	ODP 976	36.30°N, 4.31°E, 1108 m					131.9±0.9		Intensity of East Asian monsoon (grain size)	Chinese loess	35.42-37.14°N, 103.20-109.85°E	Alignment onto Holo-Stereo U-Th based chrono.				130.2±0.4 o winter [Q]		Yang and Ding (2014)
Mean ages for the beginning of ψ1-5 from Tables 3 and 4									Mean ages for the beginning of ψ1-5 from Tables 3 and 4									
136.4±1.7 133.9±0.8 133.3±1.1 130.4±1.3 128.5±1.3									136.4±1.7 133.9±0.8 133.3±1.1 130.4±1.3 128.5±1.3									
Table 3									Table 4									

Data

You will find below the data mentioned in Table 3, Table 4, Figure 8 and Figure 9 of the GMD paper

The data files have a version string (_vYYmmDD) in their file name, indicating when they have been uploaded to this site (in case we have to update them later and use a more recent version/date).

CH69-K09 (txt)
MD95-2042 (txt)
ODP976 (txt)
ODP980 (txt)
ODP983 (txt)
ODP1063 (txt)
SU90-03 (txt)
SL_LIG_Dutton2017 (txt)
d13Cstack (txt) Stack of U1308, CH69-K09 and ODP1063
IRD-stack (txt)
Figure 9 data (xlsx)

References cited

- Berger, A.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, *J. Atmospheric Sci.*, 35(12), 2362–2367, [doi:10.1175/1520-0469\(1978\)035<2362:LTVODI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2), 1978.
- Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclauss, J., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q. and Zhou, T.: PMIP4-CMIP6: the contribution of the Paleoclimate Modelling Intercomparison Project to CMIP6, *Geosci. Model Dev. Discuss.*, 1–46, [doi:10.5194/gmd-2016-106](https://doi.org/10.5194/gmd-2016-106), 2016.
- P. Koehler and C. Nehrbass-Ahles and J. Schmitt and T.F. Stocker and H. Fischer: A 156 kyr smoothed history of the atmospheric greenhouse gases CO₂, CH₄, and N₂O and their radiative forcing, *Earth System Science Data*, 9, 363–387, [doi:10.5194/essd-9-363-2017](https://doi.org/10.5194/essd-9-363-2017), 2017.
- Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B. and Wolff, E. W.: The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years, *Clim Past*, 9(4), 1733–1748, [doi:10.5194/cp-9-1733-2013](https://doi.org/10.5194/cp-9-1733-2013), 2013.

From:

<https://pmip4.lsce.ipsl.fr/> - **PMIP4**

Permanent link:

https://pmip4.lsce.ipsl.fr/doku.php/exp_design:degla_t2?rev=1549880311

Last update: **2019/02/11 11:18**

