

Contact — [Julia Hargreaves](#) to contribute to this page, or leave a comment in the discussion box.

What model runs might be relevant

There are a lot of different model runs available as part of PMIP. In fact the amount of output can be a bit overwhelming at first. One important distinction is the difference between **transient** and **equilibrium** simulations. All climate models are run forwards in time and write output every month/day. So this distinction really relates to the forcing inputs, and the climate response to them. From a practical perspective, you should be looking at the long-term statistics of the equilibrium simulations, and the temporal variations in the transient simulations. There are more details and links below, but a quick division of the experiments is...

- **PMIP4/CMIP6 Equilibrium Simulations**

- **piControl** (Preindustrial): *The control run that all the climate changes can be calculated from*
- **abrupt4xCO2** (Instantaneous quadrupling of carbon dioxide): *An idealized global warming experiment (primarily included for calculating the climate sensitivity)*
- **midHolocene** (6,000 yrs ago): *Useful for identifying impacts of orbital forcing*
- **lgm** (The last glacial maximum at 21,000 yrs ago): *The archetypal cold climate state*
- **lig127k** (Last Interglacial, 127,000 yrs ago): *Peak of the warm period prior to last glaciation*
- **Eoi400** (Pliocene, 3,205,000 years ago): *warm conditions before any Northern Hemisphere glaciation (minimal continental drift)*
- **DeepMIP** (Eocene & Paleocene, 50-60 million yrs ago): *Warm climates with carbon dioxide concentrations similar to end of century projections (continents drifted, PMIP-only)*

- **PMIP4/CMIP6 Transient Simulations**

- **1pctCO2** (Carbon dioxide concentrations increase by 1% per year): *An idealized global warming experiment (primarily included for calculating the transient climate response)*
- **historical** (1850CE onwards): *Free-running simulations of industrial era (so climate variability's phase/sign does not match observations, but its statistics should)*
- **past1000** (last millennium, 850CE onwards): *Free-running simulations seeing 1000+ years of climate forcings*
- **LDv1** (last deglaciation, 21,000 years ago to present): *Very-long simulation that encompasses all forcings (PMIP-only)*

- **Other Transient Simulations** *There are several other long palaeoclimate simulations that may be useful*

- **TraCE** This simulation runs from 22,000 years before present (22ka) to 1990 CE as well as single-forcing sensitivity simulations of varying lengths. Contact [Bette Otto-Bliesner](#)
- **MPI** Simulations with both fast and slow forcings from 8000ka-present. Contact [Johann Jungclaus](#).
- **Kiel** Simulations of both the Holocene and the Eemian exist for the Kiel model. Contact [Vyacheslav Khon](#).
- **AWI** Slowly a simulation from the Last Glacial to present is being created. Contact [Gerrit Lohmann](#)
- **IPSL** Several simulations from 6000ka to present have recently been performed in France. Contact [Pascale Braconnot](#) and Olivier Marti.
- **FAMOUS** A simulation of the last glacial cycle with accelerated-forcing. Contact [Robin Smith](#)
- **HadCM3** There are a series of equilibrium simulations of HadCM3 run over the last glacial

cycle. A version of these does exist with water isotopes. Contact [Joy Singarayer](#)

PMIP4/CMIP6 Information

Several different experiments are being performed in PMIP4.

Last Millennium

Experimental design: <http://www.geosci-model-dev-discuss.net/gmd-2016-278/>

PMIP4 website: https://pmip4.lsce.ipsl.fr/doku.php/exp_design:lm

Database: ESGF

Variables:

Mid Holocene

Experimental design: <http://www.geosci-model-dev-discuss.net/gmd-2016-279/>

PMIP4 website: https://pmip4.lsce.ipsl.fr/doku.php/exp_design:mh

Database: ESGF

Variables:

Last Glacial Maximum

Experimental design: <http://www.geosci-model-dev-discuss.net/gmd-2017-18/>

PMIP4 website: https://pmip4.lsce.ipsl.fr/doku.php/exp_design:lgm

Database: ESGF

Variables:

Last Deglaciation (21-9ka)

Experimental design: <http://www.geosci-model-dev.net/9/2563/2016/>

PMIP4 website: https://pmip4.lsce.ipsl.fr/doku.php/exp_design:index

Database: Model output mostly to be made available directly from the institutes performing the runs

Variables: ?

Eemian - Last Interglacial

Experimental design: <http://www.geosci-model-dev-discuss.net/gmd-2016-279/>

PMIP4 website: https://pmip4.lsce.ipsl.fr/doku.php/exp_design:lig127

Database: ESGF

Variables:

Pliocene

Experimental design: <http://www.clim-past.net/12/663/2016/>

USGS Website: http://geology.er.usgs.gov/egpsc/prism/7_pliomip2.html

Database: ESGF

Variables: http://geology.er.usgs.gov/egpsc/prism/7_pliomip2.html

Eocene

EECO, PETM and pre-PETM

Experimental design: <http://www.geosci-model-dev-discuss.net/gmd-2016-127/>

Website: <http://www.deepmip.org> Database: ESGF, or DeepMIP database

Variables: Outlined in Appendix A of the Experimental Design paper.

PMIP3/CMIP5 Information

PMIP Last Millenium, mid-Holocene, and LGM

The majority of PMIP model output is held in the PMIP database.

This includes model output for the three 'Tier 1' PMIP time slices: Last Millenium, mid-Holocene, and LGM.

In order to access it, contact Jean-Yves [jean-yves.peterschmitt@lsce.ipsl.fr], or see the instructions here: [database](#).

The current status of the database is here: [status](#).

PlioMIP mid-Pliocene

The PlioMIP data is held on a (password protected) ftp repository in Bristol.

This includes model output for the mid-Pliocene time slab.

In order to access the full outputs, contact Alan Haywood [a.m.haywood@leeds.ac.uk].

You can access a limited amount of output (monthly 2m air temperatures from each model, and ensemble and zonal means) from the Supplementary information of this paper:

<http://www.clim-past.net/9/191/2013/cp-9-191-2013.html>

LIGMIP last Interglaci[al/ation]

You can access a limited amount of output (monthly 2m air temperatures) from the Supplementary information of this paper: <http://www.clim-past.net/9/699/2013/cp-9-699-2013.html>

In order to access more data, you would have to contact the groups individually, or contact Dan Lunt [d.j.lunt@bristol.ac.uk] for an email list.

Bette Otto-Bliesner is currently collating precipitation data, so you could contact her [ottobli@ucar.edu] if you want that.

EoMIP Eocene

You can access a limited amount of output (monthly 2m air temperatures from each model, and ensemble and zonal means) from the Supplementary information of this paper: <http://www.clim-past.net/8/1717/2012/cp-8-1717-2012.html>

In order to access more data, you would have to contact the groups individually, or contact Dan Lunt [d.j.lunt@bristol.ac.uk] for an email list.

Other stuff

Some groups have their own online databases of model simulations:

University of Bristol BRIDGE repository:

<http://www.bridge.bris.ac.uk/resources/simulations>

If you click on 'Access simulations', you can gain access (netcdf and plots) to model data from a selection of published papers. In addition, you may be able to have access to other simulations, on request [to d.j.lunt@bristol.ac.uk or p.j.valdes@bristol.ac.uk]. A list of currently available groups of simulations is here:

http://www.paleo.bris.ac.uk/ummodel/list_of_simulations.html

[[PMIP3 Wiki Home](#)] - [[Help!](#)] - [[Wiki syntax](#)]

From:
<https://pmip4.lsce.ipsl.fr/> - **PMIP4**

Permanent link:
<https://pmip4.lsce.ipsl.fr/doku.php/wg:ptof:models?rev=1558427699>

Last update: **2019/05/21 08:34**



